



## King's Research Portal

DOI:

[10.1136/bmj.i195](https://doi.org/10.1136/bmj.i195)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

REEACT Team (2016). Corrections: Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT trial): large scale pragmatic randomised controlled trial. *BMJ (Clinical research ed.)*, 352, [i195]. <https://doi.org/10.1136/bmj.i195>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# CORRECTIONS

## Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT trial): large scale pragmatic randomised controlled trial

The original trial registration for this Research paper (*BMJ* 2015;351:h5627, doi:10.1136/bmj.h5627), the REEACT trial (<http://www.controlled-trials.com/ISRCTN91947481>), included depression as a primary outcome, measured in two ways: “Depression severity and symptomatology as measured by a validated self-report measure (the Patient Health Questionnaire [PHQ-9]) and the International Classification of Diseases (ICD-10) depression score at four months.” The original protocol defined the primary outcome measure as “depression severity and symptomatology as measured by the PHQ-9 and a self-report computer-administered diagnostic interview (CIS-R) at four months.” Before the start of the trial, the primary

outcome measure was modified to consist of only the PHQ-9 at four months. We did not collect ICD diagnosed depression using a standardised interview schedule. The public document detailing all revisions can be accessed in the full report at <http://www.journalslibrary.nihr.ac.uk/hta/volume-19/issue-101#chapters/2>. One of our outcomes was a measure of quality adjusted life years, which is published in the full economic evaluation at <http://www.journalslibrary.nihr.ac.uk/hta/volume-19/issue-101#chapters/4>.

Cite this as: *BMJ* 2016;352:i195

© BMJ Publishing Group Ltd 2016